

**ILE is indicated in refractory life-threatening cardiotoxicity secondary to local anaesthetic toxicity**

## Indications

- Cardiac arrest or life-threatening toxicity due to **local anaesthetics** in conjunction with standard ACLS
- There is currently no evidence to support the use of ILE as first line / standard treatment of life-threatening toxicity associated with other drugs
- Consider ILE as rescue therapy only in life-threatening cardiotoxicity caused by lipophilic cardio-toxins resistant to conventional treatment
- ILE is not indicated in stable dysrhythmias or seizures

## Contraindications:

- documented severe life-threatening allergy to soybean or egg

## Adverse effects:

- ECMO and dialysis filter dysfunction
- Interference with blood laboratory analysis
- Pancreatitis, acute lung injury/ARDS, fat embolism

## Presentation

- Intralipid 20% emulsion in 500mL bottle / plastic bag

## Dose and Administration

- Administer an initial bolus of 1.5 mL/kg IV over one minute
- Repeat (1.5 mL/kg IV bolus over one minute) every 5 minutes if no response to preceding dose, to a maximum of 8 doses
- Maximum cumulative total dose should NOT exceed 12 mL/kg (8 doses)
- Discontinue once haemodynamic stability achieved OR maximum dose administered

*\* If a decision is made to administer ILE as rescue therapy in life-threatening cardiotoxicity caused by lipophilic cardio-toxins resistant to conventional treatment, then discuss with a clinical toxicologist*

**Pregnancy:** Safety not established. Administration should not be withheld if indicated.