Vitamin K1 (Phytomenodione)

Vitamin K1 (Phytomenadione) is used to reverse excessive anticoagulation caused by warfarin and super-warfarin overdose.

l		
	Indications	Presentation
	Acute warfarin overdose	- 10 mg/1mL ampoule (Konakion® MM)
	- See 'Anticoagulant: warfarin' guideline	- 2 mg /0.2mL ampoule (Konakion® MM Paediatric)
	Anticoagulant rodenticides overdose	- 10 mg capsule
	- See 'Anticoagulant long-acting rodenticides'	Dose and Administration
	guideline	Patients with life-threatening haemorrhage / active uncontrolled haemorrhage / haemodynamic instability
	Contraindications:	- 10 mg Vitamin K1 intravenously (paediatric dose 0.25 mg/kg)
	- Known hypersensitivity	- 4 factor- plasma concentrate 25-50 units/kg IV via slow injection and FFP 15ml/kg IV
	Adverse effects:	Patients WITHOUT life-threatening haemorrhage / active uncontrolled haemorrhage / haemodynamic instability
	- Anaphylaxis (more common when Vitamin K1	- Vitamin K1 dosing will depend on the type of anticoagulant causing excessive anticoagulation (warfarin or
	is administered intravenously)	super-warfarins, whether the patient usually requires anti-coagulation, and whether the patient requiring
	- Facial flushing, sweating, unusual taste	anticoagulation has an absolute indication (e.g., mechanical cardiac valve or high-risk of thrombosis)
	- Venous irritation or phlebitis	See 'Anticoagulant guidelines: warfarin OR long-acting rodenticides' guidelines for specific dosing
	NOTE:	- The oral route is preferred for Vitamin K1 maintenance therapy as it is associated with less adverse effects
	- Menadione (Vitamin K3 should NOT be used	- Intravenous Vitamin K1 preparation can be given orally
	as a substitute for Vitamin K1)	Therapeutic Endpoint:
	- Vitamin K1 is not indicated for treatment of	- Stable coagulation state as clinically indicated
	excessive coagulation secondary to direct	- Stabilization may require > 48 hours, and sometimes weeks-months treatment with long-acting rodenticides
	acting oral anticoagulants (DOACs)	Pregnancy:
	e.g., dabigatran / rivaroxaban/ apixaban	- Vitamin K1 is considered safe in pregnancy

AUSTIN CLINICAL TOXICOLOGY SERVICE GUIDELINE